

# Solutionbank C1

## Edexcel Modular Mathematics for AS and A-Level

### Practice paper C1

#### Exercise 1, Question 1

**Question:**

(a) Write down the value of  $16^{\frac{1}{2}}$ . (1)

(b) Hence find the value of  $16^{\frac{3}{2}}$ . (2)

**Solution:**

(a)  $16^{\frac{1}{2}} = \sqrt{16} = 4$

(b)  $16^{\frac{3}{2}} = \left(16^{\frac{1}{2}}\right)^3 = 4^3 = 64$

© Pearson Education Ltd 2008

# Solutionbank C1

## Edexcel Modular Mathematics for AS and A-Level

### Practice paper C1 Exercise 1, Question 2

#### Question:

Find  $\int (6x^2 + \sqrt{x}) dx$ . (4)

#### Solution:

$$\begin{aligned}\int \left( 6x^2 + x^{\frac{1}{2}} \right) dx \\&= 6 \frac{x^3}{3} + \frac{x^{\frac{3}{2}}}{\frac{3}{2}} + c \\&= 2x^3 + \frac{2}{3}x^{\frac{3}{2}} + c\end{aligned}$$

© Pearson Education Ltd 2008

# Solutionbank C1

## Edexcel Modular Mathematics for AS and A-Level

### Practice paper C1 Exercise 1, Question 3

#### Question:

A sequence  $a_1, a_2, a_3, \dots, a_n$  is defined by

$$a_1 = 2, a_{n+1} = 2a_n - 1.$$

(a) Write down the value of  $a_2$  and the value of  $a_3$ . (2)

5

(b) Calculate  $\sum_{r=1}^5 a_r$ . (2)

#### Solution:

$$(a) a_2 = 2a_1 - 1 = 4 - 1 = 3$$

$$a_3 = 2a_2 - 1 = 6 - 1 = 5$$

$$(b) a_4 = 2a_3 - 1 = 10 - 1 = 9$$

$$a_5 = 2a_4 - 1 = 18 - 1 = 17$$

5

$$\sum_{r=1}^5 a_r = a_1 + a_2 + a_3 + a_4 + a_5 = 2 + 3 + 5 + 9 + 17 = 36$$

$r = 1$

# Solutionbank C1

## Edexcel Modular Mathematics for AS and A-Level

### Practice paper C1

#### Exercise 1, Question 4

**Question:**

(a) Express  $(5 + \sqrt{2})^2$  in the form  $a + b\sqrt{2}$ , where  $a$  and  $b$  are integers. (3)

(b) Hence, or otherwise, simplify  $(5 + \sqrt{2})^2 - (5 - \sqrt{2})^2$ . (2)

**Solution:**

$$(a) (5 + \sqrt{2})^2 = (5 + \sqrt{2})(5 + \sqrt{2}) = 25 + 10\sqrt{2} + 2 = 27 + 10\sqrt{2}$$

$$(b) (5 - \sqrt{2})^2 = (5 - \sqrt{2})(5 - \sqrt{2}) = 25 - 10\sqrt{2} + 2 = 27 - 10\sqrt{2}$$

$$\begin{aligned} & (5 + \sqrt{2})^2 - (5 - \sqrt{2})^2 \\ &= (27 + 10\sqrt{2}) - (27 - 10\sqrt{2}) \\ &= 27 + 10\sqrt{2} - 27 + 10\sqrt{2} \\ &= 20\sqrt{2} \end{aligned}$$

# Solutionbank C1

## Edexcel Modular Mathematics for AS and A-Level

### Practice paper C1

#### Exercise 1, Question 5

#### Question:

Solve the simultaneous equations:

$$x - 3y = 6$$

$$3xy + x = 24 \quad (7)$$

#### Solution:

$$x - 3y = 6$$

$$x = 6 + 3y$$

Substitute into  $3xy + x = 24$ :

$$3y(6 + 3y) + (6 + 3y) = 24$$

$$18y + 9y^2 + 6 + 3y = 24$$

$$9y^2 + 21y - 18 = 0$$

Divide by 3:

$$3y^2 + 7y - 6 = 0$$

$$(3y - 2)(y + 3) = 0$$

$$y = \frac{2}{3}, y = -3$$

Substitute into  $x = 6 + 3y$ :

$$y = \frac{2}{3} \Rightarrow x = 6 + 2 = 8$$

$$y = -3 \Rightarrow x = 6 - 9 = -3$$

$$x = -3, y = -3 \text{ or } x = 8, y = \frac{2}{3}$$

# Solutionbank C1

## Edexcel Modular Mathematics for AS and A-Level

### Practice paper C1

#### Exercise 1, Question 6

#### Question:

The points  $A$  and  $B$  have coordinates  $(-3, 8)$  and  $(5, 4)$  respectively.  
The straight line  $l_1$  passes through  $A$  and  $B$ .

- (a) Find an equation for  $l_1$ , giving your answer in the form  $ax + by + c = 0$ , where  $a$ ,  $b$  and  $c$  are integers. (4)
- (b) Another straight line  $l_2$  is perpendicular to  $l_1$  and passes through the origin. Find an equation for  $l_2$ . (2)
- (c) The lines  $l_1$  and  $l_2$  intersect at the point  $P$ . Use algebra to find the coordinates of  $P$ . (3)

#### Solution:

$$(a) \text{ Gradient of } l_1 = \frac{y_2 - y_1}{x_2 - x_1} = \frac{4 - 8}{5 - (-3)} = -\frac{4}{8} = -\frac{1}{2}$$

Equation for  $l_1$ :

$$y - y_1 = m(x - x_1)$$

$$y - 4 = -\frac{1}{2} \left( x - 5 \right)$$

$$y - 4 = -\frac{1}{2}x + \frac{5}{2}$$

$$\frac{1}{2}x + y - \frac{13}{2} = 0$$

$$x + 2y - 13 = 0$$

(b) For perpendicular lines,  $m_1 m_2 = -1$

$$m_1 = -\frac{1}{2}, \text{ so } m_2 = 2$$

Equation for  $l_2$  is  $y = 2x$

(c) Substitute  $y = 2x$  into  $x + 2y - 13 = 0$ :

$$x + 4x - 13 = 0$$

$$5x = 13$$

$$x = 2\frac{3}{5}$$

$$y = 2x = 5\frac{1}{5}$$

$$\text{Coordinates of } P \text{ are } \left( 2\frac{3}{5}, 5\frac{1}{5} \right)$$

# Solutionbank C1

## Edexcel Modular Mathematics for AS and A-Level

### Practice paper C1

#### Exercise 1, Question 7

#### Question:

On separate diagrams, sketch the curves with equations:

(a)  $y = \frac{2}{x}$ ,  $-2 \leq x \leq 2, x \neq 0$  (2)


(b)  $y = \frac{2}{x} - 4$ ,  $-2 \leq x \leq 2, x \neq 0$  (3)

(c)  $y = \frac{2}{x+1}$ ,  $-2 \leq x \leq 2, x \neq -1$  (3)

In each part, show clearly the coordinates of any point at which the curve meets the  $x$ -axis or the  $y$ -axis.


#### Solution:

(a)



$$y = \frac{2}{x}$$

(b) Translation of  $-4$  units parallel to the  $y$ -axis.



$$y = \frac{2}{x} - 4$$


Curve crosses the  $x$ -axis where  $y = 0$ :

$$\frac{2}{x} - 4 = 0$$

$$\frac{2}{x} = 4$$

$$x = \frac{1}{2}$$

(c) Translation of  $-1$  unit parallel to the  $x$ -axis.



$$y = \frac{2}{x+1}$$

The line  $x = -1$  is an asymptote.

Curve crosses the  $y$ -axis where  $x = 0$ :

$$y = \frac{2}{0+1} = 2$$

© Pearson Education Ltd 2008

# Solutionbank C1

## Edexcel Modular Mathematics for AS and A-Level

### Practice paper C1

#### Exercise 1, Question 8

#### Question:

In the year 2007, a car dealer sold 400 new cars. A model for future sales assumes that sales will increase by  $x$  cars per year for the next 10 years, so that  $(400 + x)$  cars are sold in 2008,  $(400 + 2x)$  cars are sold in 2009, and so on. Using this model with  $x = 30$ , calculate:

- (a) The number of cars sold in the year 2016. (2)
- (b) The total number of cars sold over the 10 years from 2007 to 2016. (3)  
The dealer wants to sell at least 6000 cars over the 10-year period.  
Using the same model:
- (c) Find the least value of  $x$  required to achieve this target. (4)

#### Solution:

(a)  $a = 400$ ,  $d = x = 30$   
 $T_{10} = a + 9d = 400 + 270 = 670$   
 670 cars sold in 2016

$$(b) S_n = \frac{1}{2}n \left[ 2a + \left( n - 1 \right) d \right]$$

So  $S_{10} = 5 [ (2 \times 400) + (9 \times 30) ] = 5 \times 1070 = 5350$   
 5350 cars sold from 2007 to 2016

(c)  $S_{10}$  required to be at least 6000:

$$\frac{1}{2}n \left[ 2a + \left( n - 1 \right) d \right] \geq 6000$$

$$5(800 + 9x) \geq 6000$$

$$4000 + 45x \geq 6000$$

$$45x \geq 2000$$

$$x \geq 44 \frac{4}{9}$$

To achieve the target,  $x = 45$ .

# Solutionbank C1

## Edexcel Modular Mathematics for AS and A-Level

### Practice paper C1

#### Exercise 1, Question 9

#### Question:

(a) Given that

$$x^2 + 4x + c = (x + a)^2 + b$$

where  $a$ ,  $b$  and  $c$  are constants:

(i) Find the value of  $a$ . (1)

(ii) Find  $b$  in terms of  $c$ . (2)

Given also that the equation  $x^2 + 4x + c = 0$  has unequal real roots:

(iii) Find the range of possible values of  $c$ . (2)

(b) Find the set of values of  $x$  for which:

(i)  $3x < 20 - x$ , (2)

(ii)  $x^2 + 4x - 21 > 0$ , (4)

(iii) both  $3x < 20 - x$  and  $x^2 + 4x - 21 > 0$ . (2)

#### Solution:

(a) (i)  $x^2 + 4x + c = (x + 2)^2 - 4 + c = (x + 2)^2 + (c - 4)$   
So  $a = 2$

(ii)  $b = c - 4$

(iii) For unequal real roots:

$$(x + 2)^2 - 4 + c = 0$$

$$(x + 2)^2 = 4 - c$$

$$4 - c > 0$$

$$c < 4$$

(b) (i)  $3x < 20 - x$

$$3x + x < 20$$

$$4x < 20$$


$$x < 5$$

(ii) Solve  $x^2 + 4x - 21 = 0$ :


$$(x + 7)(x - 3) = 0$$

$$x = -7, x = 3$$

Sketch of  $y = x^2 + 4x - 21$ :



$$x^2 + 4x - 21 > 0 \text{ when } x < -7 \text{ or } x > 3$$



Both inequalities are true when  
 $x < -7$  or  $3 < x < 5$

© Pearson Education Ltd 2008

# Solutionbank C1

## Edexcel Modular Mathematics for AS and A-Level

### Practice paper C1

#### Exercise 1, Question 10

#### Question:

(a) Show that  $\frac{(3x-4)^2}{x^2}$  may be written as  $P + \frac{Q}{x} + \frac{R}{x^2}$  where  $P$ ,  $Q$  and  $R$  are constants to be found. (3)

(b) The curve  $C$  has equation  $y = \frac{(3x-4)^2}{x^2}$ ,  $x \neq 0$ . Find the gradient of the tangent to  $C$  at the point on  $C$  where  $x = -2$ . (5)

(c) Find the equation of the normal to  $C$  at the point on  $C$  where  $x = -2$ , giving your answer in the form  $ax + by + c = 0$ , where  $a$ ,  $b$  and  $c$  are integers. (5)

#### Solution:

$$(a) (3x-4)^2 = (3x-4)(3x-4) = 9x^2 - 24x + 16$$

$$\frac{(3x-4)^2}{x^2} = \frac{9x^2 - 24x + 16}{x^2} = 9 - \frac{24}{x} + \frac{16}{x^2}$$

$$P = 9, Q = -24, R = 16$$

$$(b) y = 9 - 24x^{-1} + 16x^{-2}$$

$$\frac{dy}{dx} = 24x^{-2} - 32x^{-3}$$

$$\text{Where } x = -2, \frac{dy}{dx} = \frac{24}{(-2)^2} - \frac{32}{(-2)^3} = \frac{24}{4} + \frac{32}{8} = 10$$

Gradient of the tangent is 10.

$$(c) \text{ Where } x = -2, y = 9 - \frac{24}{(-2)} + \frac{16}{(-2)^2} = 9 + 12 + 4 = 25$$

$$\text{Gradient of the normal} = \frac{-1}{\text{Gradient of tangent}} = -\frac{1}{10}$$

The equation of the normal at  $(-2, 25)$  is

$$y - 25 = -\frac{1}{10} \left[ x - \begin{pmatrix} -2 \\ \end{pmatrix} \right]$$

Multiply by 10:

$$10y - 250 = -x - 2$$

$$x + 10y - 248 = 0$$