

# FP2 2008 Adapted

1. Solve the differential equation  $\frac{dy}{dx} - 3y = x$

to obtain  $y$  as a function of  $x$ .

(Total 5 marks)

IF  $f(x) = e^{-\int 3 dx} = e^{-3x} \Rightarrow e^{-3x} \frac{dy}{dx} - 3e^{-3x} y = xe^{-3x}$

$\Rightarrow \frac{d}{dx}(ye^{-3x}) = xe^{-3x} \Rightarrow ye^{-3x} = \int xe^{-3x} dx$

$u = x \quad v = -\frac{1}{3}e^{-3x}$   
 $u' = 1 \quad v' = e^{-3x}$

$\Rightarrow ye^{-3x} = -\frac{1}{3}xe^{-3x} + \frac{1}{3}\int e^{-3x} dx$

$\Rightarrow ye^{-3x} = -\frac{1}{3}xe^{-3x} - \frac{1}{9}e^{-3x} + c \quad \therefore y = -\frac{1}{3}x - \frac{1}{9} + ce^{3x}$


2. (a) Simplify the expression  $\frac{(x+3)(x+9)}{x-1} - (3x-5)$ , giving your answer in the form  $\frac{a(x+b)(x+c)}{x-1}$ , where  $a, b$  and  $c$  are integers. (4)

(b) Hence, or otherwise, solve the inequality  $\frac{(x+3)(x+9)}{x-1} > 3x-5$  (4)(Total 8 marks)

$\frac{(x+3)(x+9) - (3x-5)(x-1)}{(x-1)} = \frac{x^2 + 12x + 27 - 3x^2 + 8x - 5}{(x-1)}$

$\Rightarrow \frac{-2x^2 + 20x + 22}{(x-1)} = \frac{-2(x^2 - 10x - 11)}{(x-1)} = \frac{-2(x-11)(x+1)}{(x-1)}$

b)  $\frac{-2(x-11)(x+1)}{(x-1)} > 0 \Rightarrow -2(x-11)(x+1)(x-1) > 0$



$x < -1$   
 or  
 $1 < x < 11$

3. (a) Find the general solution of the differential equation

$$3 \frac{d^2 y}{dx^2} - \frac{dy}{dx} - 2y = x^2$$

(8)

- (b) Find the particular solution for which, at  $x = 0, y = 2$  and  $\frac{dy}{dx} = 3$ . (6) (Total 14 marks)

$$\begin{aligned} y &= Ae^{mx} \\ y' &= Amem^x \\ y'' &= Am^2 e^{mx} \end{aligned}$$

$$\begin{aligned} 3y'' - y' - 2y &= 0 \\ Ae^{mx}(3m^2 - m - 2) &= 0 \\ \neq 0 & \quad = 0 \end{aligned}$$

$$\begin{aligned} y &= ax^2 + bx + c \\ y' &= 2ax + b \\ y'' &= 2a \end{aligned}$$

$$\begin{aligned} (3m+2)(m-1) &= 0 \\ m = -\frac{2}{3} \quad m &= 1 \end{aligned}$$

$$y_{CF} = Ae^x + Be^{-\frac{2}{3}x}$$

$$y_{PI} = -\frac{1}{2}x^2 + \frac{1}{2}x - \frac{7}{4}$$


$$\therefore y = Ae^x + Be^{-\frac{2}{3}x} - \frac{1}{2}x^2 + \frac{1}{2}x - \frac{7}{4}$$

$$\begin{aligned} 3y'' &= 6a \\ -y' &= -b - 2ax \\ -2y &= -2c - 2bx - 2ax^2 \\ x^2 &= (6a - b - 2c) - (2a + 2b)x - 2ax^2 \end{aligned}$$

$$\begin{aligned} \therefore a = -\frac{1}{2} \quad b = \frac{1}{2} \quad -3 - \frac{1}{2} - 2c &= 0 \\ c &= -\frac{7}{4} \end{aligned}$$

4. The diagram above shows the curve  $C_1$  which has polar equation  $r = a(3 + 2 \cos \theta), 0 \leq \theta < 2\pi$  and the circle  $C_2$  with equation  $r = 4a, 0 \leq \theta < 2\pi$ , where  $a$  is a positive constant.

- (a) Find, in terms of  $a$ , the polar coordinates of the points where the curve  $C_1$  meets the circle  $C_2$ . (4)



The regions enclosed by the curves  $C_1$  and  $C_2$  overlap and this common region  $R$  is shaded in the figure.

- (b) Find, in terms of  $a$ , an exact expression for the area of the region  $R$ . (8)

- (c) In a single diagram, copy the two curves in the diagram above and also sketch the curve  $C_3$  with polar equation  $r = 2a \cos \theta, 0 \leq \theta < 2\pi$ . Show clearly the coordinates of the points of intersection of  $C_1, C_2$  and  $C_3$  with the initial line,  $\theta = 0$ . (3) (Total 15 marks)

$$a) \quad a(3 + 2 \cos \theta) = 4a \Rightarrow 2 \cos \theta = 1 \Rightarrow \cos \theta = \frac{1}{2} \quad \theta = \frac{\pi}{3}, -\frac{\pi}{3} \quad (4a, \frac{\pi}{3}); (4a, -\frac{\pi}{3})$$



$$\text{Area} = 2 \times [\text{Sector} + \text{Cardioid part}]$$

$$= 2 \left[ \frac{1}{2} (4a)^2 \frac{\pi}{3} + \frac{1}{2} a^2 \int_{\pi/3}^{\pi} (3 + 2 \cos \theta)^2 d\theta \right]$$


$$= a^2 \left[ \frac{16\pi}{3} + \int_{\pi/3}^{\pi} 9 + 12 \cos \theta + 4 \cos^2 \theta d\theta \right] \quad \begin{aligned} \cos 2\theta &= 2 \cos^2 \theta - 1 \\ 4 \cos^2 \theta &= 2 \cos 2\theta + 2 \end{aligned}$$

$$= a^2 \left[ \frac{16\pi}{3} + \int_{\pi/3}^{\pi} 11 + 12 \cos \theta + 2 \cos 2\theta d\theta \right]$$

$$= a^2 \left( \frac{16\pi}{3} + [11\theta + 12 \sin \theta + \sin 2\theta]_{\pi/3}^{\pi} \right)$$

$$= a^2 \left( \frac{16\pi}{3} + [(11\pi) - (\frac{11\pi}{3} + 6\sqrt{3} + \frac{\sqrt{3}}{2})] \right) = a^2 \left( \frac{16\pi}{3} + \frac{22\pi}{3} - \frac{13\sqrt{3}}{2} \right) = \frac{a^2}{6} (76\pi - 39\sqrt{3})$$

c)



5. (a) Find, in terms of  $k$ , the general solution of the differential equation

$$\frac{d^2x}{dt^2} + 4\frac{dx}{dt} + 3x = kt + 5, \text{ where } k \text{ is a constant and } t > 0. (7)$$

For large values of  $t$ , this general solution may be approximated by a linear function.

- (b) Given that  $k = 6$ , find the equation of this linear function. (2) (Total 9 marks)

$$\begin{aligned} x &= Ae^{mt} \\ x' &= Am e^{mt} \\ x'' &= Am^2 e^{mt} \end{aligned}$$

$$\begin{aligned} x'' + 4x' + 3x &= 0 \\ Ae^{mt}(m^2 + 4m + 3) &= 0 \\ \neq 0 \quad = 0 \\ (m+3)(m+1) &= 0 \\ m = -3 \quad m = -1 \end{aligned}$$

$$\begin{aligned} x &= at + b \\ x' &= a \\ x'' &= 0 \end{aligned}$$

$$\begin{aligned} x'' &= 0 \\ + 4x' &= 4a \\ + 3x &= 3at + 3b \\ \underline{kt + 5} &= \underline{3at + 4a + 3b} \end{aligned}$$

$$\begin{aligned} \therefore a &= \frac{1}{3}k & \frac{4}{3}k + 3b &= 5 \\ & & \Rightarrow b &= -\frac{4}{9}k + \frac{5}{3} \end{aligned}$$

$$\therefore x_{cf} = Ae^{-t} + Be^{-3t}$$

$$\Rightarrow x_{PI} = \frac{1}{3}kt - \frac{4}{9}k + \frac{5}{3}$$

$$\therefore x = Ae^{-t} + Be^{-3t} + \frac{1}{3}kt - \frac{4}{9}k + \frac{5}{3}$$

b) as  $t \rightarrow \infty$   $Ae^{-t} + Be^{-3t} \rightarrow 0$   $x \rightarrow 2t - 1$


6. (a) Find, in the simplest surd form where appropriate, the exact values of  $x$  for which

$$\frac{x}{2} + 3 = \left| \frac{4}{x} \right| \quad (5)$$

- (b) Sketch, on the same axes, the line with equation  $y = \frac{x}{2} + 3$  and the graph of

$$y = \left| \frac{4}{x} \right|, \quad x \neq 0. \quad (3)$$

- (c) Find the set of values of  $x$  for which  $\frac{x}{2} + 3 > \left| \frac{4}{x} \right|$ . (2)(Total 10 marks)



$$\frac{4}{x} = \frac{x}{2} + 3$$

$$\frac{4}{x} = -\frac{x}{2} - 3$$

$$8 = x^2 + 6x$$

$$8 = -x^2 - 6x$$

$$x^2 + 6x - 8 = 0$$

$$x^2 + 6x + 8 = 0$$

$$(x+3)^2 = 17$$

$$(x+4)(x+2) = 0$$

$$x = -3 \pm \sqrt{17}$$

$$x = -4 \quad x = -2$$

$$-4 < \frac{x}{2} < -2 \quad \text{or} \quad x > -3 + \sqrt{17}$$

7. (a) Show that the substitution  $y = vx$  transforms the differential equation

$$\frac{dy}{dx} = \frac{x}{y} + \frac{3y}{x}, \quad x > 0, \quad y > 0 \quad (I)$$

into the differential equation  $x \frac{dv}{dx} = 2v + \frac{1}{v}$ . (II) (3)

- (b) By solving differential equation (II), find a general solution of differential equation (I) in the form  $y = f(x)$ . (7)

Given that  $y = 3$  at  $x = 1$ , (c) find the particular solution of differential equation (I). (2)

$$y = vx$$

$$\frac{dy}{dx} = x \frac{dv}{dx} + v \quad \frac{d^2y}{dx^2} = x \frac{d^2v}{dx^2} + 2 \frac{dv}{dx}$$

$$\Rightarrow x \frac{dv}{dx} + v = \frac{x}{vx} + \frac{3vx}{x} \Rightarrow x \frac{dv}{dx} + v = \frac{1}{v} + 3v \Rightarrow x \frac{dv}{dx} = \frac{1}{v} + 2v$$

$$\Rightarrow x \frac{dv}{dx} = \frac{2v^2 + 1}{v} \Rightarrow \int \frac{v}{2v^2 + 1} dv = \int \frac{1}{x} dx$$

$$\Rightarrow \frac{1}{4} \ln |2v^2 + 1| = \ln |x| + C$$

$$\Rightarrow \ln |2v^2 + 1| = \ln x^4 + 4C \Rightarrow 2v^2 + 1 = Ax^4 \quad A = e^{4C}$$

$$\Rightarrow v = \sqrt{\frac{Ax^4 - 1}{2}} \Rightarrow v = \sqrt{Bx^4 - \frac{1}{2}} \quad B = \frac{A}{2}$$

$$\Rightarrow \frac{y^2}{x^2} = Bx^4 - \frac{1}{2} \Rightarrow y = \sqrt{Bx^6 - \frac{1}{2}x^2}$$

$$\text{at } (1, 3)$$

$$9 = B - \frac{1}{2} \therefore B = \frac{19}{2}$$

$$\therefore y = \sqrt{\frac{19}{2}x^6 - \frac{1}{2}x^2}$$


8. The curve  $C$  shown in the diagram above has polar equation

$$r = 4(1 - \cos\theta), \quad 0 \leq \theta \leq \frac{\pi}{2}.$$

At the point  $P$  on  $C$ , the tangent to  $C$  is parallel to the line  $\theta = \frac{\pi}{2}$ .

(a) Show that  $P$  has polar coordinates  $(2, \frac{\pi}{3})$ . (5)

The curve  $C$  meets the line  $\theta = \frac{\pi}{2}$  at the point  $A$ . The tangent at  $P$  meets the initial line at the point  $N$ . The finite region  $R$ , shown shaded in the diagram above, is bounded by the initial line, the line  $\theta = \frac{\pi}{2}$ , the arc  $AP$  of  $C$  and the line  $PN$ .



(b) Calculate the exact area of  $R$ .

(8)

parallel to  $\theta = \frac{\pi}{2} \Rightarrow \frac{dx}{d\theta} = 0$      $x = r \cos\theta = 4(1 - \cos\theta)\cos\theta$


$$\Rightarrow x = 4\cos\theta - 4\cos^2\theta$$

$$\Rightarrow \frac{dx}{d\theta} = -4\sin\theta + 8\cos\theta\sin\theta$$

$$\Rightarrow 8\cos\theta\sin\theta = 4\sin\theta \Rightarrow \cos\theta = \frac{1}{2}$$


$$\therefore \theta = \frac{\pi}{3}$$

$$r = 4(1 - \cos\theta) \quad \theta = \frac{\pi}{3} \quad r = 4(1 - \frac{1}{2}) = 2 \quad \therefore P(2, \frac{\pi}{3})$$



$$\Rightarrow ON = 1$$

$$NP = \sqrt{3}$$



$$\therefore R = \frac{\sqrt{3}}{2} + \frac{1}{2} \int_{\frac{\pi}{3}}^{\frac{\pi}{2}} 16(1 - \cos\theta)^2 d\theta$$

$$R = \frac{\sqrt{3}}{2} + 8 \int_{\frac{\pi}{3}}^{\frac{\pi}{2}} 1 - 2\cos\theta + (\frac{1}{2}\cos 2\theta + \frac{1}{2}) d\theta = \frac{\sqrt{3}}{2} + 8 \int_{\frac{\pi}{3}}^{\frac{\pi}{2}} \frac{3}{2} - 2\cos\theta + \frac{1}{2}\cos 2\theta d\theta$$

$$R = \frac{\sqrt{3}}{2} + 4 \int_{\frac{\pi}{3}}^{\frac{\pi}{2}} 3 - 4\cos\theta + \cos 2\theta d\theta = \frac{\sqrt{3}}{2} + 4 \left[ 3\theta - 4\sin\theta + \frac{1}{2}\sin 2\theta \right]_{\frac{\pi}{3}}^{\frac{\pi}{2}}$$

$$R = \frac{\sqrt{3}}{2} + 4 \left[ \left( \frac{3\pi}{2} - 4 \right) - \left( \frac{\pi}{3} - 2\sqrt{3} + \frac{\sqrt{3}}{4} \right) \right] = \frac{\sqrt{3}}{2} + 4 \left[ \frac{\pi}{2} + \frac{7\sqrt{3}}{4} - 4 \right]$$

$$R = \frac{\sqrt{3}}{2} + 2\pi + 7\sqrt{3} - 16 \quad \therefore R = 2\pi + \frac{15\sqrt{3}}{2} - 16.$$

9

$$(x^2 + 1) \frac{d^2 y}{dx^2} = 2y^2 + (1 - 2x) \frac{dy}{dx} \quad (I)$$

(a) By differentiating equation (I) with respect to  $x$ , show that

$$(x^2 + 1) \frac{d^3 y}{dx^3} = (1 - 4x) \frac{d^2 y}{dx^2} + (4y - 2) \frac{dy}{dx} \quad (3)$$

Given that  $y = 1$  and  $\frac{dy}{dx} = 1$  at  $x = 0$ ,

(b) find the series solution for  $y$ , in ascending powers of  $x$ , up to and including the term in  $x^3$ . (4)

(c) Use your series to estimate the value of  $y$  at  $x = -0.5$ , giving your answer to two decimal places. (1)

$$\frac{d}{dx} \left[ (x^2 + 1) \frac{d^2 y}{dx^2} \right] = \frac{d}{dx} (2y^2) + \frac{d}{dx} \left[ (1 - 2x) \frac{dy}{dx} \right]$$

$$= 2x \frac{d^2 y}{dx^2} + (x^2 + 1) \frac{d^3 y}{dx^3} = 4y \frac{dy}{dx} - 2 \frac{dy}{dx} + (1 - 2x) \frac{d^2 y}{dx^2}$$

$$\Rightarrow (x^2 + 1) \frac{d^3 y}{dx^3} = (1 - 4x) \frac{d^2 y}{dx^2} + (4y - 2) \frac{dy}{dx} \quad \Rightarrow$$

$$x_0 = 0 \quad y_0 = 1 \quad y'_0 = 1 \quad \Rightarrow (1) y''_0 = 2(1)^2 + (1 - 2(0)) y'_0 \Rightarrow y''_0 = 2 + 1 = 3$$

$$\Rightarrow (1) y'''_0 = (3) + 2(1) \quad \Rightarrow y'''_0 = 5$$

$$\therefore y = 1 + x + \frac{3}{2}x^2 + \frac{5}{6}x^3$$

$$x = -0.5 \Rightarrow y \approx 0.77$$

10 The point  $P$  represents a complex number  $z$  on an Argand diagram such that

$$|z - 3| = 2|z|.$$

- (a) Show that, as  $z$  varies, the locus of  $P$  is a circle, and give the coordinates of the centre and the radius of the circle. (5)

The point  $Q$  represents a complex number  $z$  on an Argand diagram such that

$$|z + 3| = |z - i\sqrt{3}|.$$

- (b) Sketch, on the same Argand diagram, the locus of  $P$  and the locus of  $Q$  as  $z$  varies. (5)
- (c) On your diagram shade the region which satisfies

$$|z - 3| \geq 2|z| \text{ and } |z + 3| \geq |z - i\sqrt{3}|. \quad (2)$$


$$|(x-3) + iy| = 2|x + iy| \Rightarrow (x-3)^2 + y^2 = 4x^2 + 4y^2$$

$$\Rightarrow x^2 - 6x + 9 + y^2 = 4x^2 + 4y^2 \Rightarrow 3x^2 + 3y^2 + 6x = 9$$

$$\Rightarrow x^2 + 2x + y^2 = 3 \Rightarrow (x+1)^2 + y^2 = 2^2$$

locus is a circle  $C(-1, 0) r=2$

$$x=0 \Rightarrow y^2=3 \Rightarrow y=\sqrt{3}$$



Q (must pass through centre)

$$|z - 3| \geq 2|z|$$

$\Rightarrow$  within circle

$$|z + 3| \geq |z - i\sqrt{3}|$$

$\Rightarrow$  closer to  $i\sqrt{3}$

11. De Moivre's theorem states that  $(\cos \theta + i \sin \theta)^n = \cos n\theta + i \sin n\theta$  for  $n \in \mathbb{R}$

(a) Use induction to prove de Moivre's theorem for  $n \in \mathbb{Z}^+$ . (5)

(b) Show that  $\cos 5\theta = 16\cos^5\theta - 20\cos^3\theta + 5\cos\theta$  (5)

(c) Hence show that  $2\cos\frac{\pi}{10}$  is a root of the equation

$$x^4 - 5x^2 + 5 = 0$$

$$n=1 \quad (\cos \theta + i \sin \theta)^1 = \cos \theta + i \sin \theta \quad \cos n\theta + i \sin n\theta = \cos \theta + i \sin \theta$$

$\therefore$  true when  $n=1$

$$\text{assume true if } n=k \quad \Rightarrow (\cos \theta + i \sin \theta)^k = \cos k\theta + i \sin k\theta$$

$$n=k+1 \quad (\cos \theta + i \sin \theta)^{k+1} = (\cos \theta + i \sin \theta)^k (\cos \theta + i \sin \theta)$$

$$= (\cos k\theta + i \sin k\theta)(\cos \theta + i \sin \theta)$$

$$= (\cos k\theta \cos \theta - \sin k\theta \sin \theta) + i(\sin k\theta \cos \theta + \cos k\theta \sin \theta)$$

$$= \cos(k\theta + \theta) + i \sin(k\theta + \theta) = \cos(k+1)\theta + i \sin(k+1)\theta$$

$$\Rightarrow (\cos \theta + i \sin \theta)^{k+1} = \cos(k+1)\theta + i \sin(k+1)\theta$$

$\therefore$  true for  $n=1$  and true for  $n=k+1$  if true for  $n=k$

$\therefore$  by Mathematical Induction true for  $n \in \mathbb{Z}^+$

$$b) (\cos \theta + i \sin \theta)^5 = \cos 5\theta + i \sin 5\theta$$

$$\begin{array}{r} 11 \\ 12 \\ 133 \\ 14641 \\ 15101051 \end{array}$$

$$(\cos \theta + i \sin \theta)^5 = \cos^5 \theta + 5i \cos^4 \theta \sin \theta - 10 \cos^3 \theta \sin^2 \theta - 10i \cos^2 \theta \sin^3 \theta + 5 \cos \theta \sin^4 \theta + i \sin^5 \theta$$

$$\text{equating real parts } \Rightarrow \cos 5\theta = \cos^5 \theta - 10 \cos^3 \theta \sin^2 \theta + 5 \cos \theta \sin^4 \theta$$

$$\therefore \cos 5\theta = \cos^5 \theta - 10 \cos^3 \theta (1 - \cos^2 \theta) + 5 \cos \theta (1 - 2 \cos^2 \theta + \cos^4 \theta)$$

$$\therefore \cos 5\theta = \cos^5 \theta - 10 \cos^3 \theta + 10 \cos^5 \theta + 5 \cos \theta - 10 \cos^3 \theta + 5 \cos^5 \theta$$

$$\therefore \cos 5\theta = 16 \cos^5 \theta - 20 \cos^3 \theta + 5 \cos \theta \quad \#$$

$$c) x = 2 \cos \theta \Rightarrow x^4 - 5x^2 + 5 = 16 \cos^4 \theta - 20 \cos^2 \theta + 5$$

$$\Rightarrow x^4 - 5x^2 + 5 = \frac{\cos 5\theta}{\cos \theta} = 0 \Rightarrow \cos 5\theta = 0 \Rightarrow 5\theta = \frac{\pi}{2}, \frac{3\pi}{2}, \dots$$

$$\therefore x = 2 \cos \frac{\pi}{10} \text{ must be a root of the equation } x^4 - 5x^2 + 5 = 0 \quad \therefore \theta = \frac{\pi}{10}, \frac{3\pi}{10}, \dots$$